arXiv:1511.03340v3 [math.DG] 21 Nov 2015

Singularities of functions with harmonic leading terms of two variables

Yuki Yasuda

November 24, 2015

Abstract

We study the classification problem of singularities of function-germs with harmonic leading terms of two variables under the right-equivalence. We study the classification in the cases that the order of function-germs is at most 7. Moreover, we observe that the multiple actions of Laplacian appear for the classifications of such class of function-germs (Theorems 1 - 3).

1 Introduction

In this paper, we study the classification of singularities of function-germs of two variables with a non-zero harmonic leading term under the right-equivalence.

We study the classification in the cases that the order of function-germs is at most 7. Moreover, we observe that the multiple actions of Laplacian appear for the classifications of such class of function-germs (see Theorems 1 - 3).

For example, we prove that singularities of functions with harmonic leading terms of order 5 are completely classified by the action of Δ^3 to the term of degree 6.

Recall that a function h on $\mathbf{R}^n$ is called a harmonic function if it satisfies the Laplace equation $\Delta h = 0$, where $\Delta = \sum_{j=1}^{n} \frac{\partial^2}{\partial x_j^2}$ denotes the Laplacian. In this paper, we study singularities of functions of two variables x and y. Therefore we put $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$.

Recall that any harmonic function-germ on $(\mathbf{R}^2, \mathbf{0})$ is a real part of a holomorphic function on $(\mathbf{C}, 0)$.

Thoughout this paper, we use the notation for the harmonic polynomials of special type:

$$f_k = \mathrm{Re}(x+iy)^k, \quad g_k = \mathrm{Im}(x+iy)^k = \mathrm{Re}(-i(x+iy)^k),$$

where $i = \sqrt{-1}$ and Re (resp. Im) means the real (resp. imaginary) part.

Let $\mathcal{H}_k(\mathbf{R}^2)$ denote the vector space of all homogeneous harmonic polynomial-germs from $(\mathbf{R}^2, \mathbf{0})$ to $(\mathbf{R}, \mathbf{0})$ of degree k. It is known that $\mathcal{H}_k(\mathbf{R}^2)$ is spanned by f_k and g_k. In particular, $\dim_{\mathbf{R}} \mathcal{H}_k(\mathbf{R}^2) = 2$.

By the following propositon, we obtain the right-classification of harmonic polynomial space $\mathcal{H}_k(\mathbf{R}^2) \setminus \{0\}$. This proposition provides the motivation to classify function-germs $h\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}, \mathbf{0})$ with harmonic leading terms with real variables under right equivalent.

Proposition 1. Let $h\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}, \mathbf{0})$ be a harmonic function-germ, and k be a natural number. Suppose that the order of h at the origin is equal to k. Then h is right-equivalent to f_k by a conformal diffeomorphism-germ $(\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}^2, \mathbf{0})$.

In particular, any element $h \in \mathcal{H}_k(\mathbf{R}^2) \setminus \{0\}$ is right equivalent to $f_k \in \mathcal{H}_k(\mathbf{R}^2) \setminus \{0\}$ by a linear conformal diffeomorphism-germ.

Proof. It is known that for every holomorphic function germ $h_i\colon (\mathbf{C}, 0) \to (\mathbf{C}, 0)(i = 1, 2)$, they are holomorphically right-equivalent if and only if $\mathrm{ord}(h_1) = \mathrm{ord}(h_2)$ (see [3]). By considering real part of h_1 and h_2, we obtain above statement. □

Note that for any harmonic function-germ h and for any linear conformal transformation φ, $h \circ \varphi$ is a harmonic function-germ.

Let $h\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}, 0)$ be a C^∞ function germ such that

$$h = h_k + R_{k+1},$$

where h_k is a non-zero homogeneous harmonic polynomial of degree k and $\mathrm{ord}(R_{k+1}) \geq k + 1$.

If $k = 1$, since h_1 is a non-zero linear function, the classification of above function-germs is obtained by implicit function theorem i.e. h is right-equivalent to x.

If $k = 2$, since h_2 is a non-degenerate quadratic form, the classification of this class is obtained by Morse lemma i.e. h is right-equivalent to $2xy$.

If $k = 3$, the singularity of h_3 is called D_4^-. It is proved that h is right equivalent to $x^2y - y^3$.

If $k = 4$, the singularity of h is of $X_{1,0}$ type (see [2]). There exists a real number a_0 such that $a_0^2 \neq 4$ and h is right equivalent to $x^4 - a_0x^2y^2 + y^4$ (in fact, $a_0 = -6$) in our case.

If degree of leading terms is equal to 5, the singularity is called of N_{16} type (see [2]). There exists a real number a, b, c such that $ab \neq 9$, $4(a^3 + b^3) + 27 - a^2b^2 - 18ab \neq 0$ and h is right equivalent to $x^4y + ax^3y^2 + bx^2y^3 + xy^4 + cx^3y^3$. Note that in [2], several invariants such as multiplicity and moduli number are studied for N_{16}.

The following result gives the complete solution to our classification problem in the case $k = 5$:

Theorem 1. (1) Any function-germ $h = h_5 + R_6\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}, 0)$ with non-zero harmonic leading term h_5 of degree 5 with $\mathrm{ord}(R_6) \geq 6$, is right equivalent to $f_5 + \tilde{R}_6$ for some function-germ $\tilde{R}_6$ with $\mathrm{ord}(\tilde{R}_6) \geq 6$ by a linear conformal diffeomorphism-germ.

(2) For any homogeneous polynomial ρ_6 of degree 6 and function-germ R_7 with $\mathrm{ord}(R_7) \geq 7$, the function-germ $f_5 + \rho_6 + R_7$ is right equivalent to $f_5 + \frac{c}{6!}x^6$ where $c = \Delta^3\rho_6 \in \mathbf{R}$.

(3) For any $c, \tilde{c} \in \mathbf{R}$, $f_5 + cx^6$ is right equivalent to $f_5 + \tilde{c}x^6$ if and only if $c = \tilde{c}$.

Other theorems (Theorem 2, Theorem 3) give pre-normal forms of function-germs in our classes for $k = 6, 7$. These theorems give us hidden relations between singularities of functions of our type and actions of Laplacian of several times.

To state Theorems 2, 3 we recall basic terms of singularity theory.

We define k-jet of h at the origin $j^k h(\mathbf{0})$ by the Taylor series of $h\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}, 0)$ considered up to the degree k.

We say that two function germs $h_i (i = 1, 2)$ are k-jet equivalent (written $h_1 \cong_{j^k} h_2$) if $j^k h_1(\mathbf{0}) = j^k h_2(\mathbf{0})$.

We say that $h_i\colon (\mathbf{R}^m \mathbf{0}) \to (\mathbf{R}^n, \mathbf{0}) (i = 1, 2)$ are $\mathcal{R}_k$ equivalent (written $h_1 \cong_{\mathcal{R}_k} h_2$) if there exists a diffeomorphism-germ $\tau\colon (\mathbf{R}^m, \mathbf{0}) \to (\mathbf{R}^m, \mathbf{0})$ and $\tilde{h}_2\colon (\mathbf{R}^m, \mathbf{0}) \to (\mathbf{R}^n, \mathbf{0})$ such that h_2 and $\tilde{h}_2$ are k-jet equivalent and h_1 is right-equivalent to $\tilde{h}_2$.

Theorem 2. (1) Any function germ $h = h_6 + R_7\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}, 0)$ with non-zero harmonic leading term h_6 of degree 6 with $\mathrm{ord}(R_7) \geq 7$ is right equivalent to $g_6 + \tilde{R}_7$ for some function-germ $\tilde{R}_7$ with $\mathrm{ord}(\tilde{R}_7) \geq 7$ by a conformal diffeomorphism-germ.

(2) For any homogeneous polynomial ρ_7 of degree 7 and function-germ R_8 with $\mathrm{ord}(R_8) \geq 8$, $g_6 + \rho_7 + R_8$ is $\mathcal{R}_7$-equivalent to $g_6 + \frac{c_1}{7!}x^7 + \frac{c_2}{6!1!}x^6 y$ where $c_1 = \frac{\partial}{\partial x}\Delta^3\rho_7 \in \mathbf{R}$, $c_2 = \frac{\partial}{\partial y}\Delta^3\rho_7 \in \mathbf{R}$.

(3) For any homogeneous polynomial ρ_7, ρ_8 of degree 7 and 8 and function-germ R_8 with $\mathrm{ord}(R_8) \geq 8$, $g_6 + \rho_7 + \rho_8$ is $\mathcal{R}_8$- equivalent to $g_6 + \rho_7 + \frac{c}{8!}x^8$ where $c = \Delta^4\rho_8 \in \mathbf{R}$.

Theorem 3. (1) Any function-germ $h = h_7 + R_8$ with non-zero harmonic leading term h_7 of degree 7 and $\mathrm{ord}(R_8) \geq 8$ is right equivalent to $g_7 + \tilde{R}_8$ for some function-germ $\tilde{R}_8$ with $\mathrm{ord}(\tilde{R}_8) \geq 8$ by a linear conformal diffeomorphism-germ.

(2) For any homogeneous polynomial ρ_8 and fucntion-germ R_9 with $\mathrm{ord}(R_9) \geq 9$ $g_7 + \rho_8$ is $\mathcal{R}_8$-equivalent to $g_7 + \frac{c_1}{8!}x^8 + \frac{c_2}{7!1!}x^7 y + \frac{c_3}{6!2!}x^6 y^2$, where homogeneous

$$c_1 = (\frac{\partial^2}{\partial x^2} - 3\frac{\partial^2}{\partial y^2})\Delta^3\rho_8 \in \mathbf{R}, \quad c_2 = \frac{\partial^2}{\partial x \partial y}\Delta^3\rho_8 \in \mathbf{R}, \quad c_3 = \frac{\partial^2}{\partial y^2}\Delta^3\rho_8 \in \mathbf{R}.$$

(3) For any homogeneous polynomial ρ_8 and ρ_9 of degree 8 and 9 and function-germ R_{10} with $\mathrm{ord}(R_{10}) \geq 10$, $g_7 + \rho_8 + \rho_9 + R_{10}$ is $\mathcal{R}_9$-equivalent to $f_7 +$

$\rho_8 + \frac{c_1}{9!}x^9 + \frac{c_2}{8!1!}x^8y$, where

$$c_1 = \frac{\partial}{\partial x}\Delta^4\rho_9 \in \mathbf{R}, \qquad c_2 = \frac{\partial}{\partial y}\Delta^4\rho_9 \in \mathbf{R}$$

(4) For any homogeneous polynomial $\rho_8, \rho_9, \rho_{10}$ of degree 8, 9 and 10, $g_7 + \rho_8 + \rho_9 + \rho_{10}$ is $\mathcal{R}_{10}$-equivalent to $g_7 + \rho_8 + \rho_9 + \frac{c}{10!}x^{10}$, where

$$c = \Delta^5\rho_{10} \in \mathbf{R}.$$

To conclude this section, we introduce a related notion and we give a Corollary of our classification results.

Definition 1 (see [1])**.** Let l be a non-zero interger. A function germ $h\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}, \mathbf{0})$ is called an *l-harmonic function-germ* of degree l if h satisfies the following equation:

$$\Delta^l h = 0.$$

Note that we use notation "homogeneous l-harmonic polynomial of degree k" for l-harmonic function and homogeneous function of degree k.

Corollary 1. Let h_5 be a non-zero harmonic polynomial of degree 5 and $\rho_6\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}, 0)$ be a homogeneous 3-harmonic polynomial of degree 6. Then $h_5 + \rho_6$ is right equivalent to h_5.

Proof. Let ρ_6 be a homogeneous 3-harmonic polynomial with degree 6 i.e. $\Delta^3\rho_6 = 0$.

By Theorem 1 (1), there exists a rotation $R_\theta\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}^2, \mathbf{0})$ and $c \in \mathbf{R}$ such that $h_5 + \rho_6$ and $f_5 + c\rho_6$ are right-equivalent.

Since rotation preserve Laplacian, $\Delta^3(c\rho_6 \circ R_\theta) = c\Delta^3(\rho_6) = 0$. Thus $h_5 + \rho_6$ is right-equivalent to f_5 by Theorem 1 (2). □

If $k = 6$ or 7, we obtain statements similar to Corollary 1:

Corollary 2. Let h_6 be a non-zero harmonic polynomial of degree 6.

1) Let ρ_7 be a homogeneous 3-harmonic function of degree 7. Then $h_6 + \rho_7$ is $\mathcal{R}_7$ equivalent to h_6.

2) Let ρ_8 be a homogeneous 4-harmonic function of degree 8. Then $h_6 + \rho_8$ is $\mathcal{R}_8$ equivalent to h_6.

Corollary 3. Let h_7 be a non-zero harmonic polynomial of degree 7.

1) Let ρ_8 be a homogeneous 3-harmonic function of degree 8. Then $h_7 + \rho_8$ is $\mathcal{R}_8$ equivalent to h_7.

2) Let ρ_9 be a homogeneous 4-harmonic function of degree 9. Then $h_7 + \rho_9$ is $\mathcal{R}_9$ equivalent to h_7.

3) Let ρ_{10} be a homogeneous 5-harmonic function of degree 10. Then $h_7 + \rho_{10}$ is $\mathcal{R}_{10}$ equivalent to h_7.

2 Proof of Theorems

To prove of Theorems 1 - 3, we prepare some additional notation and statements.

Let $\mathcal{E}_n$ denote the ring consisting of C^∞ map-germs $(\mathbf{R}^n, \mathbf{0}) \to \mathbf{R}$ and let $m_n \subset \mathcal{E}_n$ denote the ideal consisting of C^∞ map-germs $(\mathbf{R}^n, \mathbf{0}) \to (\mathbf{R}, \mathbf{0})$.

Let $h\colon (\mathbf{R}^n, \mathbf{0}) \to (\mathbf{R}, \mathbf{0})$ be a C^∞ function-germ. Let J_h denote the Jacobian ideal of h i.e. $J_h = \langle \frac{\partial h}{\partial x_i} | 1 \leq i \leq n \rangle_{\mathcal{E}_n}$.

We say that h is $\mathcal{R}_k$- determined if h is right-equivalent to some germ $\tilde{h}$ with $j^k\tilde{h}(\mathbf{0}) = j^k h(\mathbf{0})$.

We recall a sufficient condition of $\mathcal{R}_k$-determancy. For the proof, see [4].

Proposition 2 (see [4])**.** Let $h\colon (\mathbf{R}^n, \mathbf{0}) \to (\mathbf{R}, \mathbf{0})$ be a C^∞ map-germ. Suppose for a natural number k

$$m_n^k \subset m_n J_h + m_n^{k+1}$$

holds.

Then h is $\mathcal{R}_k$-determined.

Proposition 3. Any two elements in $\mathcal{H}_k(\mathbf{R}^2) \setminus \{\mathbf{0}\}$ are right equivalent by a conformal diffeomorphism-germ.

Furthermore,

$$\{\varphi\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}^2, \mathbf{0}); \varphi \in \mathrm{GL}_2(\mathbf{R}), f_k \circ \varphi = f_k\} = \langle \begin{pmatrix} \cos\frac{2\pi}{k} & -\sin\frac{2\pi}{k} \\ \sin\frac{2\pi}{k} & \cos\frac{2\pi}{k} \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \rangle.$$

Proof. Let $h_1, h_2 \in \mathcal{H}_k(\mathbf{R}^2) \setminus \{\mathbf{0}\}$. Then, by Proposition 1, there exists a linear conformal diffeomorphism-germ $\varphi_1, \varphi_2\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}^2, \mathbf{0})$ such that $h_1 \circ \varphi_1 = f_k$ and $h_2 \circ \varphi_2 = f_k$. Then, we obtain $h_2 = h_1 \circ \varphi_1 \circ \varphi_2^{-1}$.

Since $\varphi_1 \circ \varphi_2^{-1}$ is a conformal diffeomorphism-germ, first statement is proved.

Let $\varphi \in \mathrm{GL}_2(\mathbf{R})$ and suppose that $f_k \circ \varphi = f_k$. Then $\varphi(f_k^{-1}(\mathbf{0})) = f_k^{-1}(\mathbf{0})$.

Recall that $f_k = \mathrm{Re}(x+iy)^k$. Then, $f_k^{-1}(\mathbf{0})$ is k lines splitting plant to $2k$ parts for same angle.

There exists $c \in \mathbf{R}_{\geq 0}$ s.t.

$$\varphi \in c\langle \begin{pmatrix} \cos\frac{2\pi}{k} & -\sin\frac{2\pi}{k} \\ \sin\frac{2\pi}{k} & \cos\frac{2\pi}{k} \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \rangle.$$

However, $f_k(x, -y) = f_k \circ R_{\frac{2\pi}{k}} = f_k$. Therefore, we obtain the statement. □

Proposition 4. Let k be a natural number such that $k \leq 7$. Then f_k is $\mathcal{R}_{\max(k, 2k-4)}$-determined.

Proof. If $k = 1$ then $f_1 = x$. So by implicit theorem, for all $R_2 \in \mathfrak{m}_2^2$, $f_1 + R_2$ is right-equivalent to f_1.

If $k = 2$ then $f_2 = x^2 - y^2$. So by Morse Lemma, for all $R_3 \in \mathfrak{m}_2^3$, $f_2 + R_3$ is right-equivalent to f_2.

If $k = 3$ then $f_3 = 3(x^2y - xy^2)$. It is straightforward to obtain that

$$m_2^3 \subset m_2 J_{f_3} + m_2^4.$$

So, f_3 is $\mathcal{R}_3$-determined by using Proposition 2.

If $k=4$, $f_4 = x^4 - 6x^2y^2 + y^4$. It is straightforward to obtain that

$$m_2^5 \subset m_2 J_{f_4} + m_2^6.$$

Therefore by Proposition 2, f_4 is $\mathcal{R}_5$-determined.

Furthermore for any homogeneous polynomial p of degree 5, there exists a diffeomorphism germ $\varphi\colon (\mathbf{R}^2,\mathbf{0}) \to (\mathbf{R}^2,\mathbf{0})$ such that $j^5(f_4+p)(\mathbf{0}) = j^5(f_4 \circ \varphi)(\mathbf{0})$.

Thus, we have that f_4 is $\mathcal{R}_4$-determined.

For $k = 5, 6$ and 7, we can prove that g_k is $\mathcal{R}_6, \mathcal{R}_8, \mathcal{R}_{10}$-determined for the same way. □

Proof of Theorem 1. (1) By Proposition 1, there exists a linear conformal diffeomorphism-germ $\varphi\colon (\mathbf{R}^2,\mathbf{0}) \to (\mathbf{R}^2,\mathbf{0})$ such that $h_5 \circ \varphi = f_5$. Then, $(h_5+R_6)\circ\varphi = f_5 + R_6\circ\varphi$ and $\mathrm{ord}(R_6 \circ \varphi) \geq 6$. So, by taking $\tilde{R}_6$ as $R_6 \circ \varphi$, we obtain a statement.

(2) We define a local diffeomorphism germ $\varphi\colon (\mathbf{R}^2,\mathbf{0}) \to (\mathbf{R}^2,\mathbf{0})$ by

$$\begin{aligned}\varphi(x,y) = (&x + x^2(\frac{1}{5}a_6 + \frac{1}{25}a_4 + \frac{1}{25}a_2) + xy(\frac{1}{20}a_5 + \frac{1}{20}a_3 + \frac{1}{20}a_1) - y^2(\frac{a_6}{5}),\\ &y + x^2(\frac{1}{80}a_5 + \frac{1}{80}a_3 + \frac{1}{16}a_1) - xy(\frac{7}{20}a_6 + \frac{3}{50}a_4 + \frac{1}{100}a_2)\\ &\quad - y^2(\frac{1}{80}a_1 + \frac{1}{80}a_3 + \frac{1}{16}a_5)).\end{aligned}$$

.

This φ is obtained by a following way: Let us consider the local-diffeomorphism

$$\varphi(x,y) = (x + s_1x^2 + s_2xy + s_3y^2, y + s_4x^2 + s_5xy + s_6y^2).$$

Consider equations which are obtained by setting that $x^5y, x^4y^2, x^3y^3, x^2y^4, xy^5, y^6$ terms of $(f_5+\rho_6)\circ\varphi$ are equal to zero.

By solving the linear equiation of equation, we can determine $s_j (j = 1,2,3,4,5,6)$ to find φ.

Then, by straightforward computation, we obtain

$$(f_5+\rho_6)\circ\varphi \cong_{j^6} f_5 + x^6(a_0 + a_6 + \frac{a_2+a_4}{5})$$

and

$$a_0 + a_6 + \frac{a_2+a_4}{5} = \frac{1}{6!} \times (\Delta^3\rho_6).$$

(3) By Proposition 3 and the procedure to construct the above diffeomorphism φ, all diffeomorphism ψ giving the right equivalence of $f_5 + cx^6$ and $f_5 + \tilde{c}x^6$ has the following form:

$$\psi(x,y) = \varphi \circ \varphi_L(x,y),$$

where, φ_L is a composition of $e^{\frac{2\pi i}{k}}$ and $\bar{z}$ and φ is a diffeomorphism such that

$$\psi(\varphi_L(f_5 + cx^6)) \cong_{j^6} f_5 + \frac{x^6}{6!} \times \Delta^3\varphi_L(cx^6).$$

Since φ_L preserves Laplacian, we have

$$\Delta^3\varphi_L(cx^6) = \Delta^3 cx^6 = 6!c.$$

This proves Theorem 1. □

Proof of Theorem 2. (1) By Proposition 1, there exists a linear conformal diffeomorphism-germ $\varphi\colon (\mathbf{R}^2,\mathbf{0}) \to (\mathbf{R}^2,\mathbf{0})$ such that $h_6\circ\varphi = g_6$. Then, $(h_6+R_7)\circ\varphi = g_6+R_7\circ\varphi$ and $\mathrm{ord}(R_7\circ\varphi) \geq 7$. So, by taking $\tilde{R}_7$ as $R_7\circ\varphi$, we obtain a statement.

(2) Let $\rho_7 = \sum_{j=0}^{7} a_j x^{7-j}y^j$.

Define the local diffeomorphism $\varphi\colon (\mathbf{R}^2,\mathbf{0}) \to (\mathbf{R}^2,\mathbf{0})$ by

$$\begin{aligned}\varphi(x,y) = (x + x^2(&\frac{1}{48}a_3 + \frac{1}{24}a_5 + \frac{5}{16}a_7)\\ &+ xy(\frac{5}{336}a_2 + \frac{5}{168}a_4 + \frac{19}{336}a_6) - y^2(\frac{a_7}{6}),\\ y + x^2(&\frac{1}{42}a_2 + \frac{1}{70}a_4 + \frac{1}{42}a_6) - xy(\frac{1}{240}a_3 + \frac{1}{24}a_5 + \frac{19}{48}a_7)\\ &- y^2(\frac{1}{336}a_2 + \frac{1}{168}a_4 + \frac{5}{112}a_6)).\end{aligned}$$

Then, by a straightforward computation

$$(g_6+\rho_7)\circ\varphi \cong_{j^7} g_6 + x^7(a_0 + \frac{3}{35}a_4 + \frac{a_2+a_6}{7}) + x^6y(a_1 + \frac{3}{5}a_3 + a_5 + 7a_7).$$

Furthermore

$$\begin{aligned}a_0 + \frac{3}{35}a_4 + \frac{a_2+a_6}{7} &= \frac{1}{7!}\times(\frac{\partial}{\partial x}\Delta^3\rho_7),\\ a_1 + \frac{3}{5}a_3 + a_5 + 7a_7 &= \frac{7}{7!}\times(\frac{\partial}{\partial y}\Delta^3\rho_7).\end{aligned}$$

(3) Let $\rho_8 = \sum_{j=0}^{8} a_j x^{8-j}y^j$. Let us define a local diffeomorphism $\varphi\colon (\mathbf{R}^2,\mathbf{0}) \to$

$(\mathbf{R}^2, \mathbf{0})$ by

$$\begin{aligned}\varphi(x,y) = (x - x^3(&\frac{5}{768}a_1 + \frac{5}{768}a_3 + \frac{1}{256}a_5 + \frac{5}{768}a_7)\\ &+ x^2y(\frac{5}{336}a_2 + \frac{5}{168}a_4 + \frac{19}{336}a_6 + \frac{5}{12}a_8)\\ &+ xy^2(\frac{25}{768}a_1 + \frac{5}{256}a_3 + \frac{25}{768}a_5 + \frac{47}{768}a_7)\\ &- y^3(\frac{1}{6}a_8),\\ y + x^3(&\frac{1}{42}a_2 + \frac{1}{70}a_4 + \frac{1}{42}a_6 + \frac{1}{6}a_8)\\ &+ x^2y(\frac{47}{768}a_1 + \frac{25}{768}a_3 + \frac{5}{256}a_5 + +\frac{25}{768}a_7)\\ &- xy^2(\frac{5}{12}a_8 + \frac{5}{112}a_6 + \frac{1}{168}a_4 + \frac{1}{336}a_2)\\ &- y^3(\frac{5}{768}a_1 + \frac{1}{256}a_3 + \frac{5}{768}a_5 + \frac{35}{768}a_7)).\end{aligned}$$

Then, by a straightforward computation, we obtain

$$\begin{aligned}(g_6 + \rho_7 + \rho_8) \circ \varphi &\cong_{\mathcal{R}} g_6 + \rho_7 + \frac{x^8}{35}(a_0 + 5a_2 + 3a_4 + 5a_6 + 35a_8)\\ &= g_6 + \rho_7 + \frac{x^8}{8!} \times \Delta^4\rho_8.\end{aligned}$$

Theorem 2 is proved. □

Proof of Theorem 3. (1) By Proposition 1, there exists a linear conformal diffeomorphism-germ $\varphi\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}^2, \mathbf{0})$ such that $h_7 \circ \varphi = g_7$. Then, $(h_7 + R_8) \circ \varphi = g_7 + R_8 \circ \varphi$ and $\mathrm{ord}(R_8 \circ \varphi) \geq 8$. So, by taking $\tilde{R}_8$ as $R_8 \circ \varphi$, we obtain a statement.

(2) Let $\rho_8 = \sum_{j=0}^{8} a_j x^{8-j} y^j$.

Let the local diffeomorphism $\varphi\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}^2, \mathbf{0})$ be defined by

$$\begin{aligned}\varphi(x,y) := (x + x^2(&\frac{5}{448}a_3 + \frac{3}{224}a_5 + \frac{15}{448}a_7)\\ &- xy(\frac{1}{490}a_4 + \frac{3}{98}a_6 + \frac{3}{7}a_8) - y^2(\frac{3}{3136}a_3 + \frac{5}{1568}a_5 + \frac{15}{448}a_7),\\ y - x^2(&\frac{3}{245}a_4 + \frac{2}{49}a_6 + \frac{3}{7}a_8)\\ &- xy(\frac{9}{1568}a_3 + \frac{15}{784}a_5 + \frac{13}{224}a_7) + y^2(\frac{a_8}{7})).\end{aligned}$$

Then, by a direct computation,

$$\begin{aligned}(g_7 + \rho_8) \circ \varphi \cong_{j^8} g_6 &+ \frac{x^7}{7!} \times (\frac{\partial}{\partial x}\Delta^3\rho_7) + \frac{7x^6y}{7!} \times (\frac{\partial}{\partial y}\Delta^3\rho_7)\\ &+ \frac{28x^6y^2}{8!} \times (\frac{\partial^2}{\partial y^2}\Delta^3\rho_8).\end{aligned}$$

(3) Let $\rho_9 = \sum_{j=0}^{9} a_j x^{9-j} y^j$.
Let local diffeomorphism $\varphi\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}^2, \mathbf{0})$ be defined by

$$\begin{aligned}
\varphi(x,y) := (x &+ x^3(\frac{5}{448}a_3 + \frac{3}{224}a_5 + \frac{15}{448}a_7 + \frac{5}{16}a_9) \\
&+ x^2y(\frac{23}{1344}a_2 + \frac{29}{1568}a_4 + \frac{65}{3136}a_6 + \frac{17}{336}a_8) \\
&- xy^2(\frac{3}{3136}a_3 + \frac{5}{1568}a_5 + \frac{105}{3136}a_7 + \frac{51}{112}a_9) \\
&- y^3(\frac{5}{4032}a_2 + \frac{1}{672}a_4 + \frac{5}{1344}a_6 + \frac{5}{144}a_8), \\
y &+ x^3(\frac{1}{63}a_2 + \frac{1}{147}a_4 + \frac{1}{147}a_6 + \frac{1}{63}a_8) \\
&- x^2y(\frac{9}{1568}a_3 + \frac{15}{784}a_5 + \frac{13}{224}a_7 + \frac{33}{56}a_9) \\
&- xy^2(\frac{5}{672}a_2 + \frac{1}{112}a_4 + \frac{5}{224}a_6 + \frac{11}{168}a_8) + y^3(\frac{a_9}{7})).
\end{aligned}$$

Then, by a straightforward computation,

$$(g_7 + \rho_8 + \rho_9) \circ \varphi \cong_{j^9} g_7 + \rho_8 + \frac{x^9}{9!} \times (\frac{\partial}{\partial x}\Delta^4\rho_9) + \frac{9x^8y}{9!} \times (\frac{\partial}{\partial y}\Delta^4\rho_9)$$

(4) Let $\rho_{10} = \sum_{j=0}^{10} a_j x^{10-j} y^j$.
Let local diffeomorphism $\varphi\colon (\mathbf{R}^2, \mathbf{0}) \to (\mathbf{R}^2, \mathbf{0})$ be defined by

$$\begin{aligned}
\varphi(x,y) = (x &- x^4(\frac{9}{256}a_1 + \frac{1}{256}a_3 + \frac{3}{1792}a_5 + \frac{3}{1792}a_7) + \frac{1}{256}a_9 \\
&+ x^3y(\frac{23}{1344}a_2 + \frac{29}{1568}a_4 + \frac{65}{3136}a_6 + \frac{17}{336}a_8 + \frac{209}{448}a_{10}) \\
&+ x^2y^2(\frac{51}{896}a_1 + \frac{3}{128}a_3 + \frac{19}{128}a_5 + \frac{3}{128}a_7 + \frac{51}{896}a_9) \\
&- xy^3(\frac{5}{4032}a_2 + \frac{1}{672}a_4 + \frac{5}{4032}a_6 + \frac{5}{144}a_8 + \frac{209}{448}a_{10}) \\
&- y^4(\frac{1}{256}a_1 + \frac{3}{1792}a_3 + \frac{3}{1792}a_5 + \frac{1}{256}a_7 + \frac{9}{256}a_9), \\
y &+ x^4(\frac{1}{63}a_2 + \frac{1}{147}a_4 + \frac{1}{147}a_6\frac{1}{63}a_8 + \frac{1}{7}a_{10}) \\
&+ x^3y(\frac{61}{896}a_1 + \frac{3}{128}a_3 + \frac{9}{896}a_5 + \frac{9}{896}a_7 + \frac{3}{128}a_9) \\
&- x^2y^2(\frac{5}{672}a_2 + \frac{1}{112}a_4 + \frac{5}{224}a_6 + \frac{11}{168}a_8 + \frac{21}{32}a_{10}) \\
&- xy^3(\frac{3}{128}a_1 + \frac{9}{896}a_3 + \frac{9}{896}a_5 + \frac{3}{128}a_7 + \frac{61}{896}a_9) \\
&+ y^4(\frac{a_{10}}{7})).
\end{aligned}$$

Then, by a direct computation,

$$(g_7 + \rho_8 + \rho_9 + \rho_{10}) \circ \varphi \cong_{j^{10}} g_7 + \rho_8 + \rho_9 + \frac{x^{10}}{10!} \times (\Delta^5 \rho_{10}).$$

Thus, we have Theorem 3. □

References

[1] N. Aronszajn, T. M. Creese, L. J. Lipkin, "Polyharmonic Functions" Oxford Mathematical Monographs, 1983.

[2] V.I. Arnold, "Remarks on the stationary phase method and coxeter numbers", Russian Mathematical Surveys, 28 (1973), 14 - 44.

[3] J.W.Bruce, P.J.Giblin, "Curves and Singularities: A Geometrical Introduction to Singularity Theory", Cambridge University Press, 1992.

[4] J.N. Mather, "Stability of C^∞ mappings, III: Finitely determined mapgerms", Math.I.H.E.S., 35(1968), 127 - 156.